



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

COVID-19 vaccine effectiveness against the omicron (BA.2) variant in England



The omicron (B.1.1.529) variant, first detected in the UK on Nov 27, 2021, rapidly became the dominant strain, due in part to reduced vaccine effectiveness.¹ An increase in sequenced cases of the omicron sub-lineage BA.2 was observed in the week beginning on Jan 3, 2022.² BA.2 has a growth advantage over BA.1^{3,4} and has become the dominant strain in the UK at the time of writing. Neutralisation assays using monoclonal antibodies have suggested a small antigenic difference between BA.1 and BA.2, although sera from individuals with booster vaccinations neutralise both variants similarly.³

The UK COVID-19 vaccination programme has been in place since Dec 8, 2020, with primary courses of two doses of either BNT162b2 (Comirnaty, Pfizer-BioNTech), ChAdOx1-S (Vaxzevria, Oxford/AstraZeneca), or mRNA-1273 (Spikevax, Moderna). Booster vaccination with either BNT162b2 or a half dose (50 µg) of mRNA-1273 was introduced on Sept 14, 2021, to adults older than 50 years and those in risk groups, and on Nov 29, 2021, to all adults.

In this Comment, we estimate vaccine effectiveness against symptomatic disease and hospitalisation with BA.1 and BA.2 after one or two doses of BNT162b2, ChAdOx1-S, or mRNA-1273, and after booster doses of BNT162b2 or mRNA-1273 during a period of co-circulation. We used a test-negative case-control study design.⁵⁻⁸ Our analysis included all vaccines used in the UK. Vaccination status was included as an independent variable and effectiveness defined as 1 minus the odds of vaccination in cases, divided by the odds of vaccination in controls (appendix pp 1-3).

Between Jan 17 and March 31, 2022, there were 1127517 eligible tests from symptomatic individuals, of which 265820 were positive for BA.1, 246069 were positive for BA.2, and 615628 were negative (controls). The hospitalisation analysis included 15043 eligible tests, of which 1662 were positive for BA.1, 623 were positive for BA.2, and 12758 were controls (appendix pp 4-7).

There was no evidence of reduced vaccine effectiveness against symptomatic disease with BA.2 compared with BA.1 (figure; appendix p 9). 25 weeks or more after two doses, vaccine effectiveness was

14.8% (95% CI 12.9-16.7 against BA.1 and 27.8% (25.9-29.7) against BA.2. Booster immunisation increased protection after a week to 70.6% (68.9-72.2) against BA.1 and 74.0% (70.8-76.9) against BA.2, waning to 37.4% (35.8-39.0) against BA.1 and 43.7% (42.3-45.1) against BA.2 at 15 or more weeks after receiving the booster dose. Protection against hospitalisation was lower in people with BA.2 than in people with BA.1 in some periods but higher for others (figure); the CIs overlapped in most periods. After a booster dose, vaccine effectiveness against hospitalisation peaked at 90.8% (85.1-94.3) against BA.1 and 89.1% (80.5-94.0) against BA.2 before decreasing to 80.4% (75.6-84.3) and 56.5% (38.4-69.3), respectively, after 15 or more weeks. Due to the small number of individuals involved, there was greater uncertainty around the vaccine effectiveness estimates against hospitalisation than around those against symptomatic disease.

Overall, we found no reduction in vaccine effectiveness against symptomatic disease with BA.2 compared with BA.1. As previously observed, we found that vaccine effectiveness wanes over time,^{1,7,9,10} but there was no difference in the rate of decline between the two sub-lineages. Due to the small number of people with BA.2, we were not able to stratify by manufacturer in these analyses; however, we found little difference according to vaccine after a third dose in previous analyses of the omicron variant.¹ These findings are consistent with neutralisation assays,³ but there is discrepancy with a Danish household transmission study, which found that BA.2 was associated with increased susceptibility to infection for vaccinated individuals.⁴ Differences between the UK and Denmark studies might be explained by different vaccination and infection histories in the respective countries, or by methodological differences, and need further exploration.

Vaccine effectiveness against severe disease is higher and retained for longer than effectiveness against mild disease.⁹ Since the omicron variant became dominant, people are increasingly likely to have COVID-19 as an incidental finding rather than as the primary reason

Lancet Infect Dis 2022

Published Online
May 24, 2022
[https://doi.org/10.1016/S1473-3099\(22\)00309-7](https://doi.org/10.1016/S1473-3099(22)00309-7)

See Online for appendix




Figure: Vaccine effectiveness against symptomatic disease (A) and hospitalisation (B) following infection with the omicron sub-lineages BA.1 and BA.2 in adults aged 18 years and older in England

for admission.⁸ Using specific definitions to identify admissions with severe respiratory disease gives higher vaccine effectiveness estimates than using broader definitions, and these estimates probably reflect the true vaccine effectiveness against hospitalisation.⁸ Vaccine effectiveness against hospitalisation was similar for BA.1 and BA.2 after booster vaccination, although vaccine effectiveness might wane faster for BA.2 than for BA.1. The BA.2 analysis might be more susceptible to misclassification bias, because of cases incidentally hospitalised with COVID-19 due to the higher infection rates during the periods after the booster when BA.2 predominated (appendix p 4); this bias might explain the lower vaccine effectiveness estimates for BA.2 than for BA.1 in these periods.

We declare no competing interests. The UK Health Security Agency has legal permission, provided by Regulation 3 of The Health Service (Control of Patient Information) Regulations 2002, to process patient confidential information for national surveillance of communicable diseases and, as such, individual patient consent is not required to access records.

Freja C M Kirsebom, Nick Andrews, Julia Stowe, Samuel Toffa, Ruchira Sachdeva, Eileen Gallagher, Natalie Groves, Anne-Marie O'Connell, Meera Chand, Mary Ramsay, *Jamie Lopez Bernal
jamie.lopezbernal2@phe.gov.uk

UK Health Security Agency, London NW9 5EQ, UK (FCMK, NA, JS, ST, RS, EG, NG, A-MO'C, MC, MR, JLB); National Institute for Health and Care Research (NIHR) Health Protection Research Unit in Vaccines and Immunisation, London School of Hygiene & Tropical Medicine, London, UK (NA, MR, JLB); Guys and St Thomas' Hospital National Health Service Trust, London, UK (MC); NIHR Health Protection Research Unit in Respiratory Infections, Imperial College London, London, UK (JLB)

- Andrews N, Stowe J, Kirsebom F, et al. COVID-19 vaccine effectiveness against the omicron (B.1.1.529) variant. *N Engl J Med* 2022; **386**: 1532-46.
- UK Health Security Agency. SARS-CoV-2 variants of concern and variants under investigation in England: technical briefing 35. Jan 28, 2022. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1050999/Technical-Briefing-35-28January2022.pdf (accessed May 18, 2022).
- UK Health Security Agency. SARS-CoV-2 variants of concern and variants under investigation in England: technical briefing 36. Feb 11, 2022. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1054357/Technical-Briefing-36-11February2022_v2.pdf (accessed May 18, 2022).
- Lyngse FP, Kirkeby CT, Denwood M, et al. Transmission of SARS-CoV-2 Omicron VOC subvariants BA.1 and BA.2: evidence from Danish Households. *medRxiv* 2022; published online Jan 30. <https://doi.org/10.1101/2022.01.28.22270044> (preprint).
- Andrews N, Stowe J, Kirsebom F, et al. Effectiveness of COVID-19 booster vaccines against COVID-19 related symptoms, hospitalisation and death in England. *Nat Med* 2022; **28**: 831-37.
- Lopez Bernal J, Andrews N, Gower C, et al. Effectiveness of COVID-19 vaccines against the B.1.617.2 (delta) variant. *N Engl J Med* 2021; **385**: 585-94.
- Andrews N, Tessier E, Stowe J, et al. Duration of protection against mild and severe disease by COVID-19 vaccines. *N Engl J Med* 2022; **386**: 340-50.
- Stowe J, Andrews N, Kirsebom F, et al. Effectiveness of COVID-19 vaccines against omicron and delta hospitalisation: test negative case-control study. *medRxiv* 2022; published online April 1. <https://doi.org/10.1101/2022.04.01.22273281> (preprint).
- UK Health Security Agency. COVID-19 vaccine surveillance report: week 7. Feb 17, 2022. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1055620/Vaccine_surveillance_report_-_week_7.pdf (accessed May 18, 2022).
- Chemaitelly H, Tang P, Hasan MR, et al. Waning of BNT162b2 vaccine protection against SARS-CoV-2 infection in Qatar. *N Engl J Med* 2021; **385**: e83.